

Classification: Biological Sciences, Biophysics

Title: Protein threading by learning

Authors:

IKSOO CHANG^{1,2}, MAREK CIEPLAK^{1,3}, RUXANDRA I. DIMA⁴, AMOS MARITAN⁵, AND JAYANTH R. BANAVAR¹

¹Department of Physics, 104 Davey Laboratory, The Pennsylvania State University, University Park, PA 16802, USA

²Department of Physics, Pusan National University, Pusan 609-735, Korea

³Institute of Physics, Polish Academy of Science, 02-668 Warsaw, Poland

⁴Institute for Physical Science and Technology and Department of Chemistry and Biochemistry, University of Maryland, College Park, Maryland 20742, USA

⁵International School for Advanced Studies (SISSA) and Abdus Salam International Center for Theoretical Physics, Via Beirut 2-4, 34014 Trieste, Italy, and Istituto Nazionale di Fisica della Materia, Italy

Corresponding author: Jayanth R. Banavar, 104 Davey Laboratory, The Pennsylvania State University, University Park, Pennsylvania 16802, phone: 814-863-1089, FAX: 814-865-0978, email: jayanth@phys.psu.edu

pages: 14

figures: 4

tables: 1 and 2 supplementary tables

abstract: 74 words

paper: 40,435 characters

Abstract

Using techniques borrowed from statistical physics and neural networks, we determine the parameters, associated with a scoring function, that are chosen optimally to ensure complete success in threading tests in a training set of proteins. These parameters provide a quantitative measure of the propensities of amino acids to be buried or exposed and to be in a given secondary structure and are a good starting point for solving both the threading and design problems.

The principal objective of this paper is a demonstration of the viability of a framework, based on ideas from statistical physics and neural networks, for attacking the protein threading problem. Our work points to the difficulty associated with a commonly used statistical procedure for determining such parameters. We present the results of threading and design tests and present a singular value decomposition (SVD) analysis of the parameters which elucidate the interplay between degree of burial and secondary structure propensities in the folding problem.

The challenge of the protein folding problem (1-5) is to deduce the native state structure and thence the functionality of a protein from the knowledge of the sequence of amino acids. The successful completion of the human genome project has heightened interest in this problem. The information readily available as input are the sequences and native structures of a few thousand proteins (6). Given an entirely new sequence, one needs to have a sound strategy for determining its native state structure. A simpler problem, threading (7), relies on the belief that the total number of distinct folds in nature is only a few thousand (8) and attempts to match the new sequence with the best among a selection of possible native state structures. (A difficulty associated with threading is that due to steric constraints, one may not be able to mount a given sequence on a piece of a native structure of a different sequence. See, for example ref. 9) In order to assess the fit of a given sequence with a putative native state structure, one might use a coarse grained representation of the amino acids in a sequence and postulate a scoring function with a simple functional form. Perhaps the simplest such function is one which characterizes the propensities of the various types of amino acids to be in different environments:

$$S(\bar{s}, \bar{\Gamma}) = \sum_i \sum_m n(i, m) \epsilon(i, m) \quad , \quad (1)$$

where S is the score function which is a measure of the match of a sequence, $\bar{s}$, and target structure, $\bar{\Gamma}$, $n(i, m)$ is the number of amino acids of type i in the environment m and $\epsilon(i, m)$ is the score associated with it (10). For a given amino acid i , each of the $\epsilon(i, m)$'s may be shifted by the same arbitrary constant so that, without loss of generality, one may set $\sum_m \epsilon(i, m) = 0$. The advantages of such an environmental scoring function over pair-wise interactions between amino acids are its simplicity and the far greater ease of incorporating gaps in both sequence and in structure.

Our focus is on determining the score quantifying the match of a sequence to a putative native state structure, the most common approach for which utilizes statistical considerations (11-13), based on counting the number of amino acids in a given environment in the native state. Pioneering work by Bowie et al. (10) has shown that a simple statistically based approach with an environmental score leads to excellent results for the inverse folding problem.

Our studies used a training set of 387 proteins (see Table I in Supplementary Information) from the PDBselect (6,14) consisting of sequences varying in length from 44 to 1017 with low sequence homology and covering many different 3D-folds according to the SCOP classification (15). Additional criteria used in selecting the proteins in the training set were: a) the protein structure was obtained through X-ray crystallography, b) the structures were monomeric, c) the determined structures missed no more than two amino acids. The same criteria were used to obtain a test set of 213 distinct proteins (Table 2 in Supplementary Information) with lengths ranging between 54 and 869. For each structure, we used a simple environmental classification which consists of the local secondary structure (α -helix, β -strand or other) and the exposed area evaluated as the ratio between the accessible area of each amino acid, X, of its native sequence (having this structure as its native state) and the

corresponding area in a Gly-X-Gly extended structure. The values of the exposed area were divided into three categories of small, medium and large exposures corresponding to $< 10\%$, $10 - 50\%$, and $> 50\%$ respectively. Thus the scoring function consists of nine parameters for each amino acid corresponding to each of the nine environments that it might be found in.

Materials and Methods

We begin by applying the ideas of Bowie et al. (10) to the threading problem. The statistical score $\epsilon_s(i, m)$ associated with amino acid i in an environment m is readily deduced using the expression

$$\epsilon_s(i, m) = -\ln [P(i, m)/P(i)] \quad , \quad (2)$$

where $P(i, m)$ is the probability of finding an amino acid of type i in the environment of type m and $P(i)$ is the probability of finding an amino acid of type i in any environment. Both $P(i, m)$ and $P(i)$ are determined from a knowledge of the sequences and native state structures of the proteins in our training set. In order to assess the quality of the extracted scores, we carried out threading tests on all but the largest protein in the training set itself. Each protein sequence was mounted on its own native state structure and on every fragment (of the correct length chosen without insertions and deletions) of all the larger proteins. The exposed area for the amino acid mounted on a fragment was assumed to be the same as that in the whole protein from which the fragment was extracted. As we shall see later, this may be a poor approximation when the size of the fragment is much smaller than the whole protein. In each case, Eqn. (1) was used to determine the scoring function. While the technique is simple, the results of gapless threading tests are only moderate – the native state structure is correctly recognized for 69% of the proteins. In a recent paper, Baud and Karlin (16) considered 418 proteins and determined the frequencies of occurrence of the twenty amino

acids in nine environments which were defined in a way similar to ours. We have converted their frequencies into statistical scores (which turn out to be similar to the statistical scores derived from our training set), using equation (2), and find 54 failures in our set of 213 proteins. This moderate performance may be due to the fact that the form of the scoring function is too simple. Support for this comes from earlier work which has pointed out the difficulty of determining the optimal interactions that stabilize the native state of even 1 protein (crambin) with a more complex scoring function involving 210 pairwise interactions (17). An alternative possibility, that the statistical approach is flawed (18) would be of more serious concern because such statistical schemes are commonly used in the protein folding problem.

We turn to a demonstration that an alternative strategy based on ideas originating in statistical physics and neural networks provides a powerful framework for tackling the threading problem. Following the pioneering work of Friedrichs and Wolynes (19) and especially Goldstein et al. (20), the basic idea is to postulate the form of a scoring function and to choose its parameters to ensure that the true native states of proteins with known structures (learning set) correspond to better (lower) scores than when the sequences are housed in competing decoy conformations (17-28). An important advantage of this procedure is that it can be used to verify whether the chosen form of the scoring function is equal to the task or not. Indeed, one may start with the simplest form of the scoring function and systematically expand the parameter space until the optimal interactions are learned. The statistical procedure considers proteins and their native state structures, whereas the learning procedure has information on competing structures as well. Our scheme is similar in spirit to that of previous work with the important differences that we consider an environmental scoring function instead of a pairwise contact potential and we optimize the energy gap without any

normalization.

The total number of inequalities (one obtains the inequalities for each sequence in the training data set by considering as decoys all pieces of the native state structures of longer proteins in the training set) is over 13 million making the problem technically difficult. For a given protein, each decoy leads to a linear inequality of the form $\sum_{i=1}^{20} \sum_{m=1}^9 [n(i, m)^D - n(i, m)] \epsilon(i, m) > 0$, where $n(i, m)^D$ is the number of amino acids of type i found in the environment m in the given decoy. The perceptron procedure is a simple technique based on neural networks for simultaneously solving a set of such linear inequalities (29). We used this procedure to optimally choose the 180 parameters in order to ensure that the worst inequality (among the more than 13 millions) was satisfied as well as possible and that threading tests on the training set were 100% successful.

Results

We describe the results of several tests and a biological interpretation of these parameters:

Learning procedure versus statistical approach:

Figure 1 shows a plot of the parameters determined using the statistical approach versus those deduced by the learning procedure. The poor correlation is consistent with the qualitatively different performance levels in threading. It underscores the fundamental difficulties of the statistical approach and points to the advantage of learning the optimized parameters in a systematically expanded parameter set.

Threading tests:

The couplings $\epsilon_{387}(i, m)$ obtained based on learning the native states of the 387 proteins

in the training set were subjected to threading on the test set containing 213 distinct proteins (Table II in Supplementary Information) and the decoys obtained from their native state structures. In contrast to the performance of the statistical parameters, for which one is unable to correctly recognize the native states of 76 of the 213 proteins, the number of failures when one uses the learned parameters is 23. The failed proteins are modest in size and have sequence lengths ranging between 54 and 131. The ranks of the native states, defined as the number of better performing decoys, of the failed proteins are plotted in Figure 2 as a function of the sequence length for both sets of parameters (note the dramatically different scales of the y -axis). For the poorest performer, using the perceptron based method, there are 102 decoys (out of 37617) that perform better than the native state (protein 1abq of length 56) while the corresponding number for the statistically derived parameters is 29424 (out of 31436 decoys for protein 1vqb of length 86). We have checked that around half of the failures are spurious for the case of the learned parameters and arise because the exposed areas for the winning decoy, which is a piece of the native state structure of a longer protein, is quite different from that determined for the whole protein. This effect of an inaccurate assignment of the exposed area is strong only for small sized proteins. The remaining failures (a total of 5 %) is likely due to the identification of genuine competitors to the native state or because the winning decoy is not a viable structure for the sequence under consideration (9).

We also tested the ϵ_{387} parameters on all 600 proteins (Table I and II in Supplementary Information) and the decoys obtained using all 600 native state structures. There were 57 failures whereas the statistically derived parameters resulted in 209 failures. We used the perceptron procedure (29) to learn the scoring parameters in order to ensure that the native state of all the proteins in the training and test set were recognized with 100 % success and

the energies of all decoys were pushed up as much as possible compared to the native state energies. In the rest of the paper, we will use this refined set of optimal parameters $\epsilon(i, m)$ (Table I) to carry out our further studies. Note that the sum of the first nine entries of each row in Table I is equal to zero and the sum of the squares of all such 180 entries has been chosen to be 180.

The native state of crambin (1crn) which was not part of the training set, is recognized in threading. This result is encouraging because of earlier difficulties in learning pairwise parameters for this protein (19). It should be noted, however, that a single amino acid mutation of 1crn, the protein 1cbn, was present in the basic learning set of 387 proteins.

As a further test, we selected 26 Globin proteins from the RCSB website (<http://www.rcsb.org/pdb/>) which were in the DEOXY form, which were not mutated and whose structures are resolved well. Strikingly, 23 of the 26 proteins correctly picked their own native state from among the millions of decoy conformations obtained from the fragments of the 600 proteins in the training and test sets described previously. For the 3 other cases, fragments from the Globin family were picked to be the best structure. Indeed, the scores of the Globin proteins on fragments of other Globin structures were generally lower than on fragments of structures of unrelated proteins underscoring the quality of our scoring function.

Biological interpretation of learned parameters:

Let us begin with a geometrical picture of $\epsilon(i, m)$, considered to be twenty vectors of nine components each. For a given amino acid i , the components of the nine-dimensional vector, labelled by the index m , capture the propensities of that amino acid to be in each of nine environments. Each environment may be thought of as representing an axis in an orthogonal

9-dimensional space. Singular value decomposition (28, 31) affords a simple prescription for dimensional reduction by the optimal choice of a new set of orthogonal axes. In this new reference frame, the original vectors span a lower-dimensional space and the axes may be conveniently rank-ordered in importance.

The SVD theorem (31) states that the 20×9 (non-square) matrix ϵ can be written as

$$\epsilon = YV^T, \quad (3)$$

where Y is a 20×9 dimensional matrix and V is a 9×9 dimensional matrix. The superscript T denotes the transpose matrix. The matrix Y is given by $Y = U\Sigma$, where Σ is a 20×9 dimensional matrix whose elements are all zero except for the diagonal terms, $\Sigma_{n,n}$, $n=1,\dots,9$. These diagonal terms are equal to the square roots, σ_n , of the common eigenvalues of $\epsilon\epsilon^T$ and $\epsilon^T\epsilon$. The σ_n 's are called singular values and are assumed to be rank ordered so that σ_1 is the largest. Here, they are: 10.59, 5.02, 3.98, 3.42, 2.57, 2.09, 1.77, 0.95, and 0.0. The columns of V , denoted by V_k , are the eigenvectors corresponding to the rank ordered eigenvalues of the matrix $\epsilon^T\epsilon$ and the columns of the 20×20 matrix U , denoted as U_k , $k=1,\dots,20$, are determined by the formula $U_k = \frac{1}{\sigma_k}\epsilon V_k$ (when the singular values are non-zero; the other cases are irrelevant for the reconstruction of the ϵ parameters). The result of the SVD transformation is that $\epsilon(i, m)$ may now be represented as a sum of contributions that diminish in an overall sense as one considers smaller singular values. The n 'th contribution is given by $y_{(n)}(i) v_{(n)}^T(m)$, where the first factor depends on the amino acid and the second on the environmental index. Thus $v_{(n)}^T$ are the new orthogonal and normalized directions — or modes — in the space of environments.

Figure 3 shows the three most dominant contributions, corresponding to the top three singular eigenvalues. Each contribution is displayed in two panels. The upper panels show

the mode as a function of the nine environmental parameters. The lower panels show the corresponding amplitudes $y_{(n)}$ plotted so that $y_{(n)}$ increases monotonically.

The first mode is dominant for 13 amino acids: C, F, I, V, H, S, T, N, P, Q, E, R, and K. The second mode is the leader for W, M, Y, and D and the third for L and A. The remaining amino acid, G, is dominated by the fifth mode. The first mode provides the overall dominant behavior and strongly distinguishes between the buried and exposed environments in a monotonic way regardless of the secondary structure — $y_{(1)}$ allows one to arrange the amino acids into buried and exposed groups depending on whether it is large and positive or large and negative. One may further subdivide the two basic groups of buried (B) and exposed (E) amino acid into subgroups: B_1 , B_2 , B_3 , E_1 , and E_2 . The division is illustrated in Figure 3 and corresponds to occurrences of more rapid variations in $y_{(1)}$ as one moves from one amino acid to the next. The key point is that the amino acids in B_1 have a strong tendency to be buried and the charged amino acid K in E_2 has a strong tendency to be exposed, and most of the amino acids are more sensitive to the degree of burial than to other considerations. This tendency for burial is usually associated with hydrophobicity in the protein folding problem (32-34). The hydrophobic amino acids F, I, V, L, and A do belong to group B but this group also contains polar amino acids. Cysteine, C, shows the strongest propensity to be buried. It should be noted that a pair of C's may form a strong contact by establishing a disulfide bridge. Of the 896 C-C contacts generated in our study of 600 proteins, 402 had both C's buried whereas only in four cases were both of the C's exposed (independent of the secondary structure). This tendency alone yields a high statistical score for C being buried. (Note that 37% of the structural sites of the 600 proteins are classified as buried, 40 % as medium, and 33 % as exposed). The learned score is even further accentuated because most of the decoys correspond to C being not buried

and stability of the native state with respect to decoys is enhance by such an adjustment.

The remaining modes break the symmetry between the secondary structures. The second mode is neutral to α and favors (disfavors) β (loop) when the coefficient $y_{(2)}$ is negative. It shows a strong preference for amino acids, such as W, with a large negative $y_{(2)}$ to be in a β -strand with a large exposed area and for amino acids, such as D, with a large positive $y_{(2)}$ to be in loops with a large exposed area. The third mode introduces a preference for C, F, K, etc. to be in β -strands with medium exposure and for L, P, and A, etc. to stay either in exposed β -strands or in buried loops and avoid exposed helices.

Protein design: We turn now to an extension of our studies to protein design or the inverse folding problem. In analogy with equilibrium statistical mechanics, the probability that a sequence $\bar{s}$ is housed in a structure $\bar{\Gamma}$ is given by (35-38)

$$P_{\bar{\Gamma}}(\bar{s}) = \frac{e^{-S(\bar{s}, \bar{\Gamma})/T}}{\sum_{\Gamma'} e^{-S(\bar{s}, \Gamma')/T}} \equiv \frac{e^{-S(\bar{s}, \bar{\Gamma})/T}}{e^{-F(\bar{s})/T}} \quad (4)$$

where T , here, is a fictitious temperature, the score S has been assumed to play a role analogous to the energy and F , the free score, plays the role of the free energy. The key point is that in the limit of $T \rightarrow 0$ and when $\bar{\Gamma}$ is the native state structure of $\bar{s}$, $P \rightarrow 1$. In this limit, therefore, the "free score" which is a function of the sequence alone approaches the score of the sequence in its native state. The last column of Table 1 shows the average contribution to the native state scores, S_i , from each type of amino acid in the various environments. It is defined by $S_i = \frac{1}{N_i} \sum_{k=1}^{N_i} \epsilon(i, m(k))$, where the sum is over the N_i occurrences of amino acid i in the native state of all 600 proteins in the training and test sets. The zero "temperature" free score of a sequence may then be readily deduced without any knowledge of the structure, by adding up these contributions for the amino acids in the sequence. Figure 4 shows a plot of the native state score versus the sequence free score

for all 600 proteins. The latter, which has no structure dependent information, provides a reasonable approximation to the actual native state score. We have verified that both are linearly proportional to the protein length and for the longer proteins, the native state score is somewhat higher than the free score due to the increasing tendency toward frustration as the sequence length increases. For design purposes, the free score provides a measure of the score one is entitled to expect in a typical native state structure and the lower the score in the target native state structure with reference to the free score, the better is the design.

Stability of cold shock proteins: We used the learned parameters to provide a molecular interpretation of the different thermal stabilities of a pair of cold shock proteins (39), one of which is mesophilic *Bacillus subtilis* (Bs-CspB: 1csp) and the other thermophilic *Bacillus caldolyticus* (Bc-Csp: 1c9o). The former has a score of -34.80 in the native state, whereas the latter is more stable with a score of -41.64. More strikingly, the free scores are -28.64 and -27.97 respectively underscoring the much better design of the thermophilic protein. We also used the conformation space of all decoys to estimate the "heat capacity" of the two proteins as a function of temperature. The heat capacity which is a measure of the fluctuations in the score (viewed as an energy) shows a peak as a function of the temperature in both cases. The peak temperature, which is a measure of the folding transition temperature of the protein, of 1c9o is higher than that of 1csp, and reflects the better thermal stability of 1c9o in accord with the experimentally observed behavior (39).

Conclusion

In summary, we have shown that a straightforward learning scheme leads to the determination of excellent environmental parameters which can be used in simple threading tests. Our results point to the danger of employing statistical procedures for estimating these

values. The learned parameters capture information on the environments in the competing structures in addition to that in the native state structures and allows for a stabilization of the native state with respect to decoy structures. Our procedure validates the notion that in the simplest cases we have studied here, a simple environmental scoring function is sufficient for capturing the essential features of protein threading. Our method has the distinct advantage of ease of expanding the parameter space and opens up the possibility of using the scoring parameters determined here as a starting point for learning the penalty parameters characterizing insertion and deletion.

This work was supported by grants from NASA, INFN and MURST (Italy), the Donors of the Petroleum Research Fund administered by the American Chemical Society, PNU research fund and KBN (grant number 2P03B-146-18).

1. Anfinsen, C. (1973) *Science* **181**, 223-230.
2. Wolynes, P. G., Onuchic, J. N., & Thirumalai, D. (1995) *Science* **267**, 1619-1620.
3. Dill, K. A. & Chan, H. S. (1997) *Nature Struct. Biol.* **4** 10-19.
4. Fersht, A. P. (1998) *Structure and mechanism in protein science: A guide to enzyme catalysis and protein folding*, New York, Freeman.
5. Baker, D. A. (2000) *Nature* **405**, 39-42.
6. Berman, H. M., Westbrook, J., Feng, Z., Gilliland, G., Bhat, T. N., Weissig, H., Shindyalov, I. N. & Bourne, P.E. (2000) *Nucl. Acid. Res.* **28**, 235-242.
7. Jones, D. T., Taylor, W. R., & Thornton, J. M. (1992) *Nature (London)* **358**, 86-89.
8. Chothia, C. (1992) *Nature* **357**, 543-544.
9. Ramachandran, G. N. & Sasisekharan, V. (1968) *Adv. Prot. Chem.* **28**, 283-437.
10. Bowie, J., Lüthy, R. & Eisenberg, D. (1991) *Science* **253**, 164-170.
11. Tanaka, S. & Scheraga, H. A. (1976) *Macromolecules* **9**, 945-950.
12. S. Miyazawa, S. & Jernigan, R. L. (1985) *Macromolecules* **18**, 534-552.
13. Zhang, C. & Kim, S. (2000) *Proc. Natl. Acad. Sci.* **97**, 2550-2555.
14. Hobohm, U. & Sander, C. (1994) *Prot. Sci.* **3**, 522-524.
15. Murzin, A. G., Brenner, S. E., Hubbard, T. & Chothia, C. (1995) *J. Mol. Biol.* **247**, 536-540.

16. Baud, F. & Karlin, S. (1999) *Proc. Natl. Acad. Sci.* **96**, 12494-12499.
17. Vendruscolo, M., Najmanovich, R. & Domany, E. (1999) *Phys. Rev. Lett.* **82**, 656-659.
18. Dima, R. I., Banavar, J. R. & Maritan, A. (2000) *Protein Sci.* **9**, 812-819.
19. Friedrichs, M. S. & Wolynes, P. G. (1989) *Science* **246**, 371-373.
20. Goldstein, R., Luthey-Schulten, Z. A. & Wolynes, P.G. (1992) *Proc. Natl. Acad. Sci.* **89**, 9029-9033.
21. Koretke K.K., Luthey-Schulten, Z. A. & Wolynes, P.G. (1996) *Protein Science* **5**, 1043-1059.
22. Mayorov, V. N. & Crippen, G. M. (1992) *J. Mol. Biol.* **227**, 876-888.
23. Mirny, L. A. & Shakhnovich, E. I. (1996) *J. Mol. Biol.* **264**, 1164-1179.
24. Clementi, C. Maritan, A. & Banavar, J. R. (1998) *Phys. Rev. Lett.* **81**, 3287-3290.
25. Dima, R. I., Settanni, G., Micheletti, C. Banavar, J. R. & Maritan A. (2000) *J. Chem. Phys.* **112**, 9151-9166.
26. Vendruscolo M., Mirny L.A., Shakhnovich E. I. & Domany E. (2000) *Proteins: Structure, Function, and Genetics* **41**, 192-201.
27. Tobi, D. & Elber, R. (2000) *Proteins: Structure, Function, and Genetics* **41**, 40-46.
28. Tobi, D., Shafran, G., Linial, N. & Elber, R. (2000) *Proteins: Structure, Function, and Genetics* **40**, 71-85.
29. Krauth, W. & Mezard, M. (1987) *J. Phys. A* **20**, L745-L752.

- 30. Park, B. & Levitt, M. (1996) *J. Mol. Biol.* **258**, 367-392.
- 31. Watkins, D. S. (1991) *Fundamentals of Matrix Computations*, Wiley, New York.
- 32. Kauzmann, W. (1959) *Adv. Protein Chem.* **14**, 1-63.
- 33. Dill, K.A. (1990) *Biochemistry* **29**, 7133-7155.
- 34. Kamtekar, S., Schiffer, J. M., Xiong, H. Y., Babik, J. M. & Hecht, M. H. (1993) *Science* **262**, 1680-1685.
- 35. Deutsch, J. M. & Kurosky, T. (1996) *Phys. Rev. Lett.* **76**, 323-326.
- 36. Seno, F., Vendruscolo, M., Maritan, A. & Banavar, J. R. (1996) *Phys. Rev. Lett.* **77**, 1901-1904.
- 37. Dima, R. I., Banavar, J. R., Cieplak, M. & Maritan, A. (1999) *Proc. Natl. Acad. Sci.* **96**, 4904-4907.
- 38. Micheletti, C., Maritan, A. & Banavar, J. R. (1999) *J. Chem. Phys.* **110**, 9730-9738.
- 39. Perl, D., Mueller, U., Heinemann, U. & Schmid, F. (2000) *Nature Struct. Biol.* **7**, 380-383.

Table Captions

Table 1: Table of ϵ , the nine environmental scores for each amino acid. Large negative values indicate a strong preference for the particular environment whereas large positive values indicate an aversion. The last column shows S_i which is a measure of the average contribution of each amino acid to the native state score and provides an estimate of the expectation of the contribution of a given amino acid to the native state score.

Table I. 180 Environmental Scores

		α			β			Other			S_i
Amino Acid		Small	Med.	Expo.	Small	Med.	Expo.	Small	Med.	Expo.	
CYS	(C)	-1.29	0.07	1.81	-1.78	-0.83	3.63	-1.24	-0.85	0.49	-1.06
PHE	(F)	-0.90	-0.35	2.33	-1.77	-1.02	1.51	-0.26	-0.28	0.74	-0.73
TRP	(W)	0.41	0.32	1.64	-1.18	-1.00	-1.02	0.57	0.50	0.91	-0.07
ILE	(I)	-0.50	-0.27	0.38	-0.25	-0.39	0.61	-1.05	0.56	0.92	-0.29
VAL	(V)	0.42	0.06	-0.12	-1.48	-0.64	0.89	-0.28	0.57	0.58	-0.31
MET	(M)	-0.26	-0.36	0.65	-0.52	0.71	1.26	-0.24	-0.77	-0.47	-0.30
LEU	(L)	-0.33	-0.16	0.09	-0.32	0.83	-0.76	-0.54	0.77	0.41	-0.10
GLY	(G)	0.36	1.16	0.73	-0.05	0.16	0.14	-0.49	-0.95	-1.06	-0.48
TYR	(Y)	0.13	0.83	-0.06	-0.42	-1.18	-0.23	0.23	0.08	0.63	0.00
ALA	(A)	-0.40	-0.05	-0.13	0.27	0.50	-0.15	-0.23	0.35	-0.25	-0.06
HIS	(H)	1.05	-0.60	-0.82	0.62	0.56	0.14	-0.29	-0.08	-0.57	-0.09
ASP	(D)	-0.29	-0.79	-0.90	1.31	0.93	1.32	0.59	-0.99	-1.17	-0.60
SER	(S)	-0.31	-0.01	-0.98	0.48	0.78	-0.75	1.00	-0.32	-0.10	0.03
THR	(T)	0.80	0.49	-0.46	0.55	-0.50	-0.80	0.74	-0.34	-0.48	0.01
ASN	(N)	0.67	-0.66	-0.66	1.34	0.60	-0.06	0.55	-0.48	-1.30	-0.39
PRO	(P)	2.35	-0.28	-0.88	1.32	1.03	-0.30	-1.02	-0.62	-1.61	-0.65
GLN	(Q)	1.74	-0.84	-1.24	0.94	-0.87	-1.07	1.32	0.01	0.01	-0.26
GLU	(E)	0.83	-0.81	-1.28	1.67	-0.21	-0.67	1.60	0.04	-1.16	-0.53

ARG	(R)	2.29	-0.80	-1.37	1.37	-1.16	-1.35	1.82	0.13	-0.94	-0.38
LYS	(K)	1.20	-1.13	-1.77	4.32	-1.43	-1.91	2.38	-0.32	-1.35	-1.11

Supplementary Table I. PDB codes for 387 learning proteins

1CII 1KCW 4HB1 1DHX 1BLE 1SQC 1AB4 1PKP 1FTY 1RGS 1TDJ 1FSZ 8LDH 6ICD
 1AEP 1ANV 5PTD 1A0I 1B6E 1SIG 1DIV 1KXU 1BVB 1BAJ 1A6F 1LRV 2ITG 1CBY
 1LXA 1914 1CC5 1A0P 1OHK 1JON 1PJR 1A7J 1AUA 1GLN 1IHP 1HLB 1ZAP 2STV
 1A17 1RDR 1RLW 1UBY 2SAS 1BCO 1A8Y 1AX8 2LIV 1AN8 2OMF 1A41 1C25 1AK5
 1AQT 1AJ6 2FXB 1BOB 1INP 1CYX 1XSM 1BIA 1CPT 1OPR 1PLQ 1AUQ 1KIT 1CTN
 1DHR 1OBR 1RCB 1A26 1CIY 1GPC 1PFO 1GRJ 1BY9 1MAZ 1LBA 1KTE 1AM2 1BB9
 1HTP 1BIX 1TUL 1DRW 1AQE 2GSQ 1DHS 5EAU 1CFR 1GEN 1BR9 1ACC 1YGS 1B5L
 1IAM 1A32 1RMD 1PEA 1SEK 1KLO 1OXA 1CRB 2TCT 1ESC 1TFR 2NG1 1LCI 1PHT
 1ALY 1VIN 1A6Q 1A76 1A1X 1CSN 1TIG 1A8H 1BTN 1CDY 1CFB 1MSC 1AMX 1HOE
 1UOX 2PGD 1BV1 2PLC 4MT2 1SRA 1DDT 1NSJ 1UOK 1POC 1SUR 1GOX 1GSA 1MJC
 2PIA 1LKI 1BY2 1SKF 1BIF 1PBV 1ALO 1RMG 2I1B 1DPE 1AJ2 4PAH 1FCE 1PNE
 1BF2 1AZ9 1A53 3TDT 7TAA 1OPC 1PTQ 1BEA 1PUC 1FUA 1RSS 1ECL 1SKZ 1NEU
 1ALU 1CUK 1CA1 1MAI 1AD2 1OPY 1EDT 1BHE 1JDW 1PHM 1DXY 1VOM 1CEO 1A8L
 1TMY 1SVB 1AIL 1WHO 1JDC 1SFP 3TSS 1DUN 1HOW 1PBE 1GPR 1A48 4ENL 2PII
 3GCB 1BG7 1VLS 1PUD 2ABK 1MDL 1RKD 1EUR 1DMR 1GND 1UCH 1BG2 1AK0 1UXY
 2GAR 1LCL 1MML 1POT 1QNF 1NPK 1AYL 1TIF 1BD8 1BDO 1BG6 1C3D 1HYP 2POR
 1UAE 1BJ7 1TML 1TYV 1HCL 2SAK 1FNA 1AL3 2TGI 2ACY 1LST 1LBU 1AMP 1NAR
 1FAS 2CBP 1FMB 1AXN 1TUD 1PDA 1HA1 1CV8 1CHD 1AMF 1USH 1CPQ 1BM8 1XWL
 1BGC 1AJJ 1TFE 1NKR 1IDO 1VJS 1BHP 1WAB 1VIE 1VHH 1GCA 1PDO 1FDR 1PMI
 1SBP 1GOF 1AKO 1MOF 2GDM 1FXD 1FNC 1GAI 2HFT 1OSA 1VNS 3CHY 1ERV 1DHN
 1AQB 1CNV 119L 1CEM 1CXC 1VCC 1GVP 2DRI 1MBA 1A3C 1EDG 1PHF 16PK 451C
 1B6A 1BKF 1RZL 5NUL 1AOP 1A8E 1CVL 1ARV 1NIF 3CYR 1MRJ 1ZIN 1LAM 1CSH
 1KUH 1PTF 1BFG 1BFD 3PTE 2AYH 2MYR 1NOX 1AKR 2A0B 1A8D 1MOQ 1HFC 1RA9
 1TCA 3GRS 2CBA 1KPF 5ICB 1AIE 1KOE 1WHI 1RIE 1MLA 1HKA 1OPD 1FLP 2MCM
 1CYO 1POA 1BRT 2HBG 2SNS 1XNB 2RN2 3SEB 1BGF 2END 1YGE 3VUB 2CTC 1HMT
 1PPT 1BQK 1UTG 1PLC 1BK0 1DCS 1C52 7RSA 1OAA 1MSI 1YCC 2PTH 2SN3 1AMM
 1BX7 1ATG 2KNT 1MUN 1A7S 1CTJ 1BS9 2IGD 1NKD 3SIL 2ERL 1A6M 1CEX 1IXH
 1BYI 1AHO 1NLS 2FDN 3LZT 1RB9 3PYP 1CBN 1GCI

Supplementary Table II. PDB codes for 213 testing proteins

1AF5 1FHE 1CRY 1ADT 2CHR 2LDX 2UCZ 1AVC 1LDB 1AR2 1BMP 1ABQ 1TLK 1ULA
 1EIA 1A1S 1B5M 1PEX 3PHV 1OJT 2FHI 6FIT 1A43 2TPT 1YFM 1LU1 1CPY 1HJP
 1ILE 1FBL 1GWZ 1BIK 1PMT 1A06 4FXC 1BAG 1HUP 1CYW 1JSG 1BMG 1FSU 2CND
 2GPR 1PBK 2ABL 1TMO 1CYG 2ALR 1NAT 1SBF 2TDX 1KAS 1A6I 4P2P 1SZT 1DOL
 1FGS 1AC5 1HIB 1QPG 1CQA 1DOT 1MKP 1ANN 8OHM 1ASS 1JNK 1BET 8CHO 2ASR
 1AD6 1BQG 1DHY 1A45 1DIK 1CIU 3KAR 1APA 2PK4 1HEY 1BYT 1ODD 1ZXQ 1FTS
 1AA0 1GDD 1PHK 1VIP 1HAR 1TSY 1AW9 1ENY 1AVK 1YVS 1BFS 5PNT 1A0K 1AIR
 2DAP 1ECY 2ASI 1ANU 1CBG 1FAJ 1P38 1A6O 1BCG 1A6L 3ERK 1BED 1BKL 1ENH
 1BLU 1A80 1A8Z 1KIV 1MNC 1A3K 1XAA 1A0B 2PSR 1B56 1LPP 1BK2 3KVT 1NFO
 1INR 1RGP 1AQ1 1TN3 2E2C 1GSH 1A8P 1AZ5 1PVL 1BKM 1VPE 1MHO 1BK1 1AE7
 1FIL 1MN1 1RCI 1RIS 1HVF 1ACF 1BDB 1AYI 1NHP 1IAG 1CGT 1AA2 2EBN 1BK9
 1POH 1ESL 1HCZ 1A58 1AEW 1BAM 1SNP 1AK2 3GAR 1EIF 1CLC 4RHN 1DFX 1A8B
 1KVY 1CYI 1UKZ 1DOI 1BGP 1TN4 1CYJ 1YMV 1AX0 1HFX 1LRA 1RCY 1NUC 1JUG
 3PRN 1ENP 1LML 1DYR 1OBM 1A44 1ZRN 2HTS 1RFS 1AMK 1IVD 1IFT 1NFN 1ARS
 1AHR 1XND 1RDS 1A68 1VQB 1ALQ 1MRG 1A7E 2SLI 1GBS 3KIV 1BZA 1AT5 2ERA
 1A3D 1GZI 1A8S

Figure Caption

Figure 1: Plot of the optimal ϵ parameters versus those determined using a statistical scheme, ϵ_s , using a training set of 387 proteins.

Figure 2: Results of the threading tests for 213 proteins arranged according to their length, N . Only the failed cases are shown. The top panel shows a plot of the number of decoys that performed better than the native state structure versus N whereas the bottom panel shows a similar plot for the couplings that were determined statistically. Note the disparity in the scales of the y -axes.

Figure 3: The top three contributions to $\epsilon(i, m)$ as emerging from the SVD analysis. The numbers in the ovals indicate the mode number. The letters at the top left of each segment of two panels indicate amino acids (in the single letter code) for which this particular mode is dominant. The top panels in each segment show the modes – the values of $v_{(n)}^T$ for the nine values of the environmental variable m . For each kind of secondary structure, the environments are listed in order from the small to large exposure. The bottom panels show the amino-acid-dependent weights $y_{(n)}$ with which the displayed mode contributes to the score in a given environment.

Figure 4: Plot of the zero "temperature" free score and the native state score of each of the proteins in the training and test sets.

FIGURES




FIG. 1.




FIG. 2.






FIG. 3.




FIG. 4.